

The changes of vitamin C, chlorophyll, carotene, xanthophylls content and the antiradicalic activity of sunflower sprouts after selenium treatment

Camelia Moldovan¹, Nicoleta Hădărugă¹, Diana Raba¹, Mirela Popa¹,
Aurica Borozan², Mărioara Drugă¹, Delia Dumbravă¹

¹Faculty of Agrofood Processing Technology, Banat's University of Agricultural Sciences and Veterinary Medicine "King Michael I of Romania" from Timișoara

²Faculty of Horticulture and Forestry, Banat's University of Agricultural Sciences and Veterinary Medicine "King Michael I of Romania" from Timișoara

Received: 02 March 2017; Accepted: 24 March 2017

Abstract

In this paper we present some of the consequences of the selenium treatment on sunflower sprout. We determined: vitamin C, chlorophyll pigments, xanthophylls and carotenes content and antiradical activity (AA). Sunflower sprouts were obtained by sunflower seeds germination in presence of sodium selenite (5 and 10 ppm Se) – experimental group, and water – control group. The methods used in our study were: Vitamin C – by iodometric method, AA – by DPPH method, chlorophyll pigments, xanthophylls and carotenes – by spectrophotometric method. The selenium treatment has led to increasing all of these parameters in experimental group, comparing to control group. The selenium treatment of sunflower seeds improves the nutritive quality of sunflower sprouts with beneficial effect on the human consumption.

Keywords: sunflower sprout, selenium, vitamin C, chlorophyll, carotene, xanthophylls, antiradicalic activity.

1. Introduction

The nutritionists of our day are focused on healthy nutrition. Fruits of vegetables are unanimously considered healthy by them [13], but the sprouts (germinated seeds and shoots) are some of the most valuable from nutritional point of view. It is known that there are major differences regarding the nutritional value of seed compared to the sprout. By germination, functional foods can be obtained [15], so sprouts provide very high nutrient levels [1, 2, 15, 17, 18]. The sprouts are rich in phytochemical nutrients, vitamins, minerals, enzymes and amino acids, having great importance in human health [13, 15]. For its wealth of nutrients and its high conversion rate of anorganic compounds in organic compounds [12], sprouts are suitable for selenium fortified, especially in European countries, where

selenium deficiency was observed and different strategies are followed to fortify foods [7]. There are numerous studies [4, 7, 9, 10, 14, 15, 18, 22, 23] where selenium supplementation of different sprouts (sunflower, rucola, wheat, barley, oats, lettuce, green tea, clover, rape, lupine, garden cress). The consequences of selenium supplementation were observed about different biochemical parameters like antioxidant activity [3, 7, 9, 18, 21], chlorophyll's pigments [10, 11, 23], carotenoid content [10, 11, 22, 23], vitamin C [4, 14, 20]. Thus, the researchers have reported that selenium supplementation determines an increasing of chlorophyll's pigments on rucola, on wheat [11, 23], lettuce [10]. The variation of carotenoids level, consecutive to selenium treatment was reported in wheat [10, 23], rucola [11], carrot [22].

The level of vitamin C increased significantly in green tea [20], lettuce [19], wheat [4, 14], clover, sunflower [14], barley and oats [4]. Antiradical activity improved after treatment with selenium of rape [9], lupine [8], clover [3, 21], garden cress [7], sunflower [18].

In this paper we present some of the consequences of the selenium treatment on sunflower sprout. We determined: vitamin C, antiradical activity (AA), chlorophyll pigments, xanthophylls and carotens.

2. Materials and Methods

Sunflower sprouts were obtained by sunflower seeds germination on textile supports. There were two experimental groups whom it was administered sodium selenite: E1(5 ppm Se) and E2 (10 ppm Se), and a control group (C) which we have given water. All of sprout samples were given in the same environmental conditions (temperature, humidity, light). Our experiment was conducted over a period of 14 days, after which was evaluated the content of: chlorophylls, xanthophylls and carotens, vitamin C and antiradical activity.




Figure 1. Scheme of assessing the level of key antioxidants in the sunflower seedlings consecutive sodium selenite treatment

The chlorophyll pigments, xanthophylls and carotenes level were assessed by spectrophotometric method using UV-VIS Perkin Elmer - Lambda 25 spectrophotometer and vitamin C - by iodometric method. All of these methods were presented by [5, 6, 16].

3. Results and Discussion

The selenium treatment has led to increasing all of these parameters in experimental group, comparing to control group.

The results of the chlorophyll pigments level of sunflower sprout analyzed samples are shown in Figure 2. From the total chlorophyll, the chlorophyll "a" has share of majeure, while the chlorophyll "b" has insignificant share.

The experimental groups with added selenium doubled their total chlorophylls and chlorophyll "a" level (18.056, respectively 21.38 mg/g) compared to control group (10.882 mg/g). The increases of chlorophyll level were directly correlated with selenium doses. This fact was similar with other research on rucola [11], on wheat [16, 23], and lettuce [10].

The results about level of the carotenes and xanthophylls in sunflower sprout analyzed samples are shown in Figure 3. The carotenes and xanthophylls level was substantially increased (double or even triple content) following selenium treatment of the sunflower sprout,

The increase in carotenoids content, consecutive to selenium treatment was reported in wheat [10, 16, 23], rucola [11] and carrot [22].




Figure 2. Chlorophyll pigments level in sunflower sprouts after selenium treatment (mg/g)

Carotens and xanthophylls (microg/g)




Figure 3. Carotens and xanthophylls level in sunflower sprouts after selenium treatment ($\mu\text{g/g}$)

In the Figure 4 are present the results of the vitamin C level from sunflower sprout analyzed samples. Our results have shown beneficial effects of selenium supplementation on sunflower sprout. In the experimental groups was registered a higher (3.048, respectively 4.931 mg/100 g) content of vitamin C, compared to control group (0.194 mg/100 g). The literature data presents similar aspects regarding vitamin C – which increased significantly after selenium treatment in green tea [20], lettuce [19], wheat [4, 14, 16], clover, sunflower, alfaalfa [14], barley and oats [4,16].

The antiradical activity of sunflower sprout samples are shown in Figure 5. As expected (all of mentioned parameters until now are known as antioxidants) antiradical activity of sunflower sprout was significantly improved in selenium exposed groups, showing direct correlation with chlorophylls, vitamin C, carotens and xanthophylls.

There are a lot of other research works which reported that antiradical activity was improved after selenium treatment in rape [9], lupine [8], clover [3, 21], garden cress [7], sunflower [18].




Figure 4. Vitamin C level in sunflower sprouts after selenium treatment (mg/100g)

Antiradicalic activity (microM/s)




Figure 5. Antiradicalic activity in sunflower sprouts after selenium treatment ($\mu\text{M/g}$)

Conclusion

The selenium treatment of sunflower seeds improves the nutritive quality of sunflower sprouts with beneficial effect on the human consumption.

- The selenium treatment (5 respectively 10 ppm Se, as sodium selenite) of sunflower sprouts had beneficial effects on antioxidant contents: chlorophyll pigments, vitamin C, carotens and xanthophylls, which lead to overall improvement of antioxidant activity.
- The increasing of antioxidants level in sunflower sprout was direct correlated with the increasing of selenium dose (until 10 ppm Se).

The nutritional value of sunflower sprouts increased consecutive to selenium treatment.

Compliance with Ethics Requirements. Authors declare that they respect the journal's ethics requirements. Authors declare that they have no conflict of interest and all procedures involving human / or animal subjects (if exist) respect the specific regulation and standards.

References

1. Balasaraswathi R., Sadasivam S., Changes in oil, sugars and nitrogenous components during germination of sunflower seeds, *Helianthus annuus*, *Plant Foods Hum Nutr.*, **1997**, 51(71). 71–77, <https://doi.org/10.1023/A:1007924026633>.
2. Cuddeford D., 1989. *Hydroponic grass*, In Practice. **1989**, 11(5), 211–214, <http://dx.doi.org/10.1136/inpract.11.5.211>.
3. de la Luz Mora María, Pinilla Ligia, Rosas Analí, Cartes Paula, Selenium uptake and its influence on the antioxidative system of white clover as affected by lime and phosphorus fertilization, *Plant and Soil, Biomedical and Life Sciences*, 303(1-2), 139-149, **2008**, Springer Netherlands, <https://doi.org/10.1007/s11104-007-9494-z>.
4. Dîma M., *The qualitative evaluation of grain fortified with selenium*. Doctoral thesis, Jelgava: LLU, **2010**, 50–95.
5. Dumbravă Delia-Gabriela, Moldovan Camelia, Raba Diana-Nicoleta, Popa Mirela-Viorica, Comparative analysis of vitamin C content and antioxidant activity of some fruits extracts, *Journal of Agroalimentary Processes and Technologies*, **2012**, 18(3), 223-228.
6. Dumbravă Delia-Gabriela, Moldovan Camelia, Raba Diana-Nicoleta, Popa Mirela-Viorica, Vitamin C, chlorophylls, carotenoids and xanthophylls content in some basil (*Ocimum basilicum* L.) and rosemary (*Rosmarinus officinalis* L.) leaves extracts, *Journal of Agroalimentary Processes and Technologies*, **2012**, 18(3), 253-258.
7. Frias Juana, Gulewicz P, Martinez-Villaluenga C, Peñas E, Piskula MK, Kozłowska H, Ciska E, Gulewicz K, Vidal-Valverde C., Changes in Nutritional Value and Cytotoxicity of Garden Cress Germinated with Different Selenium Solutions, *Journal of Agricultural and Food Chemistry*, **2010**, 58(4), 2331-2336. DOI:10.1021/jf903458u
8. Frias Juana, Gulewicz P., Martinez-Villaluenga Cristina, Pilarski R., Blazquez E., Jimenez B., Gulewicz K., Vidal-Valverde C., Influence of Germination with Different Selenium Solutions on Nutritional Value and Cytotoxicity of Lupin Seeds, *J. Agric. Food Chem.*, **2009**, 57(4), 1319–1325, DOI: [10.1021/jf8028368](https://doi.org/10.1021/jf8028368).
9. Hasanuzzaman M., Hossain M.A., Fujita M., Selenium in higher plants: Physiological role, antioxidant metabolism and abiotic stress tolerance. *J. Plant Sci.*, **2010**, 354-375, DOI: [10.3923/jps.2010.354.375](https://doi.org/10.3923/jps.2010.354.375).
10. Hawrylak-Nowak Barbara, Effect of selenium on selected macronutrients in maize plants, *J. Elementol.* **2008**, 13(4), 513-519.
11. Khattab H., Metabolic and oxidative responses associated with exposure of *Eruca sativa* (rocket) plants to different levels of selenium. *Inter J of Agricul Biol.* **2004**; 6:1010–1106.
12. Kopsell D.A., Kopsell D.E., Randle W.M. Seed Germination Response of Rapid-Cycling Brassica oleracea Grown Under Increasing Sodium Selenate, *Journal of Plant Nutrition*, **2003**, 26(7), 1355-1366.
13. Krinsky Norman I., Johnson Elizabeth J., Carotenoid actions and their relation to health and disease, *Molecular Aspects of Medicine*, **2005**, 26(6), 459-516, DOI: [10.1016/j.mam.2005.10.001](https://doi.org/10.1016/j.mam.2005.10.001).
14. Lintschinger, L., N. Fuchs, J. Moser, D. Kuehnelt, Goessler W., Selenium-Enriched Sprouts. A Raw Material for Fortified Cereal-Based Diets, *Journal of Agricultural and Food Chemistry* **2000**, 48(11), 5362–5368.
15. Marton M., Mandoki Zs., Zs. Csapo-Kiss, Csapo J., *The role of sprouts in human nutrition*. A review. *Acta Univ. Sapientiae, Alimentaria*, 3, **2010**, 81-117.
16. Moldovan Camelia, Hădăruță Nicoleta, Dumbravă Delia, Raba Diana, Popa Mirela, Toța Cristina, Assessing the level of key antioxidants in wheat seedlings consecutive sodium selenite treatment, *Journal of Agroalimentary Processes and Technologies*, **2011**, 17(1), 58-64.
17. Omi Laila, Imtiyaz Murtaza, Seed sprouting: a way to health promoting treasure, *IJCRR*. **2014**, 6(23), 70-74.
18. Pařak Paulina, Socha R., Gałkowska Dorota, Roźnowski J., Phenolic profile and antioxidant activity in selected seeds and sprouts, *Food Chemistry* **2014**, 143, 300-306, DOI: [10.1016/j.foodchem.2013.07.064](https://doi.org/10.1016/j.foodchem.2013.07.064).
19. Qingmao S., Gao L., Li S., Effect of Selenium on quality of hydroponics lettuce, *Journal of China Agricultural University*, **1998**.
20. Xu J., Fangmei Yang, Licheng Chen, Yun Hu, Qiuhui Hu, Effect of selenium on increasing the antioxidant activity of tea leaves harvested during the early spring tea producing season, *J Agric Food Chem.* 51(4), **2003**, 1081-1084, DOI: [10.1021/jf020940y](https://doi.org/10.1021/jf020940y).
21. Zielinska Danuta, Frias Juana, Piskula M. Konrad, Kozłowska Halina, Zielinski H., Vidal-Valverde C., Evaluation of the antioxidant capacity of lupin sprouts germinated in the presence of selenium, *Eur Food Res Technol.* **2008**, 227, 1711–1720, <https://doi.org/10.1007/s00217-008-0898-7>.
22. Wang D., Grieve C.M., Suarez D.L., Composition of Irrigation Water Salinity Affects Growth Characteristics and Uptake of Selenium and Salt Ions by Soybean, *Journal of Plant Nutrition, Volume*, **2005**, 28(6), 1073-1088, DOI: [10.1081/PLN-200058909](https://doi.org/10.1081/PLN-200058909).
23. Yao Xiaoqin, Chu Jianzhou, Wang Guangyin, Effects of Selenium on Wheat Seedlings Under Drought Stress, *Humana Press Inc.*, 283-290, **2008**, <https://doi.org/10.1007/s12011-009-8328-7>.