

Lignin-degrading Bacteria

Ghellai Lotfi

¹Laboratory of Applied Microbiology in Food, Biomedical and Environment (LAMAABE), Department of Biology, University of Tlemcen, 13000 Tlemcen, Algeria

Received: 11 March 2014; Accepted: 30 March 2014

Abstract

Degradation of lignin by microorganisms was primarily highlighted in rot and brown-rot fungi, which are capable of producing certain extracellular lignolytic enzymes. Recently, studies on bacteria have also shown that these organisms likely possess a set of extracellular oxidative enzymes implicated in lignin degradation. Therefore, from a biotechnological point of view, lignolytic bacteria could have several novel applications regarding the valorisation and modification of the lignin biopolymers.

Keywords: lignin, lignolytic bacteria, lignases

1. Introduction

Lignin, a polymer composed of phenylpropanoid units, associated with cellulose and hemicellulose in plant cell, is the most abundant aromatic substance present in the biosphere. Instead, it is composed of a number of chemically distinct subunits, or monolignols, the abundance of which can vary among species, among individuals, and even among cell types within an organism [1]. The decay of lignin in nature is a complex phenomenon not yet completely understood. The ability of lignin to resist degradation can be attributed to its distinctive polymeric structure [2] and its complicated aromatic polymer and high molecular weight make it resistant to microorganisms, resulting in the prevention of effective cellulose utilization [3]. Indeed, complete degradation of lignin is mainly assured by microorganisms. However, only a few bacterial species are currently known to degrade lignin molecule. Furthermore, microbial degradation of lignin has been well studied in white-rot (*Phanerochaete chrysosporium* and *Trametes versicolor*) and

brown-rot fungi (*Fomitopsis palustris*), but is much less well studied in bacteria [4].

Among these white rot fungi, basidiomycete, *P. chrysosporium* has become the most commonly used organism due to its ability to produce ligninolytic enzymes, fast growth and easy handling during culturing techniques [5]. However, there are some constraints that should be overcome by lignolytic organisms for attacking and decomposing the lignin. Fungi are capable of overcoming these constraints by their extracellular lignolytic enzymes called ligninases, including heme-dependent lignin peroxidases, manganese peroxidases, versatile peroxidases [6,7,8], and multicopper-dependent laccases [9].

White-rot fungi produce a range of extracellular lignolytic enzymes. The first molecular information on bacterial high molecular weight metabolite formation was reported with the isolation of a secreted bacterial heme peroxidase from a gram-positive bacterium, *Streptomyces viridosporus* T7A [10, 11].

Thus, there are indications that bacteria use similar types of extracellular lignin-degrading enzymes to fungi [4]. But in general, the heme peroxidases from studied lignolytic bacteria have been found to be less oxidatively powerful compared to the fungal enzymes involved in lignin degradation [12].

However, it is possible that the major lignolytic enzymes in bacterial systems is not the heme peroxidases. Beyond the enzymes used by fungi, bacteria also contain a suite of oxidative enzymes that could modify lignin for breakdown by hydroxylation or demethylation, such as cytochrome P450s, non-heme iron enzymes, or Mn- and Cu-containing oxidases [12]. As indicated before, owing of the physical properties of lignin and its incorporation into the cell wall, current approaches used for its removal from biomass are sufficiently expensive and energy intensive [13]. Although certain lignocellulolytic fungi are able to secrete industrial quantities of extracellular enzymes, bacterial enzyme production can be more cost-efficient [14]. This is because they grow more rapidly, produce multi-enzyme complexes with increased functionality and higher specificity, and can tolerate larger and more diverse environmental stress [15,16]. Lignocellulolytic bacteria could also potentially allow better separation of lignin from cellulose and thereby increase the value of both lignin, which is currently a waste product, and cellulose [14]. Moreover, It has been shown that fungi despite of there high capability of lignin-degrading, they are sometimes not stable in practical treatment under extreme environmental and substrate conditions [17]. In contrast, bacteria, in particular, deserve to be studied for ligninolytic potential because of their immense environmental adaptability and biochemical versatility. This article will focus on the diversity of lignolytic bacteria.

2. Lignolytic bactreria

Insolubility of lignin and its lack of stereoregularity contribute to making it a substrate that is difficult for the microflora to degrade [18]. Furthermore, the need to gain access to the substrate by penetration of plant tissues constitute an additional challenge for microorganisms, bacteria in particular. Actinomycetes and fungi,

accomplish this task by hyphal invasion of the various cell wall layers [18]. Nonfilamentous bacteria, originally thought to be unable to fulfill this requirement, have been found to erode wood fibers through both tunneling [19,20] and cavitation [21]. In contrast, information available indicates that lignin is fairly recalcitrant to bacterial attack, at least under laboratory conditions [18]. For this reason, some authors believe that bacteria play a secondary role in lignin biodegradation in natural environments [22,23]. However, bacteria seem to play a leading role in decomposing lignin in aquatic ecosystems [18]. Both fungi and bacteria have been observed to metabolize lignin; however, their differential reactivity with this substrate indicates that they may utilize different chemical strategies for its breakdown [12]. Although the metabolism of lignin is not as complete compared to fungal systems, it is clear that bacteria can react with lignin and possibly produce smaller aromatics that can be imported into the cell for aromatic catabolism, which is also widespread in soil bacteria [24, 25]. Bacterial lignin degradation mainly occurs under aerobic conditions. Lignin-degrading activity under aerobic conditions has been reported from diverse groups of bacteria [26]. Indeed, bacterial systems have been found to be less oxidatively powerful compared to lignolytic fungal systems to date [12]. The few bacterial species currently known to degrade cellulose and lignin are within *Pseudomonas* (order Pseudomonadales), *Cellulomonas* (order Actinomycetales), *Streptomyces* (order Actinomycetales), and other genera within the order Actinomycetales [27, 28] and are likely employing extracellular laccases and peroxidases [14]. Many studies Suggest that *Pseudomonas* sp. is the most efficient lignin degradation bacterium [29]. It not only has the ability to degrade nature lignin but it also degrades aromatic rings [30]. Thus, bacteria of other genera, including *Alcaligenes*, *Arthrobacter*, and *Nocardia*, readily degrade the single-ring aromatic compounds that build up the lignin macromolecule [31]. Clostridiales was previously regarded as a degrader of cellulose and hemicellulose, but in recent years, it was also reported participating in lignin degradation [32]. There are several reports of bacteria isolated from termite guts that have aromatic degradation capability [33] such as *Rhodococcus erythropolis* [34], *Microbacterium* sp., *Brucella melitensis*,

Ochrobactrum sp. and *Sphingomonas* sp. [35], *Burkholderia cepacia* [36], *Burkholderia* sp. and *Burkholderia* sp. [37]. However, there are conflicting reports on the ability of termite gut micro-organisms to break down lignin [4]. furthermore, bacteria isolated from wood-boring beetles possess in vitro lignolytic activity such as *Paracoccus* spp., *Escherichia* spp., *Yersinia* spp., *Rahnella* spp., *Pantoea* and *Erwinia* [38, 39, 40]. Recently a bacteria consortium containing at

least nine genera (*Clostridiales*, *Geovibrio thiophilus*, *Desulfomicrobium*, *Pseudomonas* sp., *Azoarcus* sp., *Thauera*, *Paenibacillus* sp., *Cohnella* sp., *Acinetobacter* sp., *Microbacterium*, and uncultured bacterium) was found mainly responsible for lignin degradation, based on various screening procedures [41]. Some other bacteria are known to date, to have activity for lignin breakdown are listed in Table 1.

Table 1. Lignolytic bacteria and their substrate-degrading

Bacteria species	Substrate-degrading	References
<i>Streptomyces viridosporus</i> T7A	depolymerize lignin	[42]
<i>Pseudomonas paucimobilis</i> SYK-6	break down various dimeric lignin compounds	[43]
<i>Pseudomonas putida</i> mt-2 <i>Rhodococcus jostii</i> RHA1	Lignin degradation	[44]
<i>Azotobacter</i> sp. HM121	lignin mineralization and solubilization	[45]
<i>Aneurinibacillus aneurinilyticus</i>	depolymerize lignin	[46]
<i>Bacillus</i> sp.	degrade 37% kraft lignin by culture at 30 °C for 6 days	[47]
<i>Paenibacillus</i> sp	degrade kraft lignin	[48]

3. Conclusion

According to several previous studies, Lignin degradation which is a complex process with synergism among more enzymes, is mainly assumed by both bacteria and fungus. Many bacterial strains can solubilize and metabolize lignin compounds but their ability to mineralize polymeric lignin is limited compared to fungi. Furthermore, it was found that bacterial consortiums were more effective in lignocelluloses degradation than other isolates [3]. Recent studies on lignin degradation in nature may provide novel resources for the delignification of dedicated bioenergy crops and other sources of lignocellulosic biomass [2]. Indeed, Lignocellulolytic bacteria have promised to be a fruitful source of new enzymes for next-generation lignocellulosic biofuel production [14].

Compliance with Ethics Requirements: Authors declare that they respect the journal's ethics requirements. Authors declare that they have no conflict of interest and all procedures involving human and/or animal subjects (if exists) respect the specific regulations and standards.

References

- Ralph, J.; Lundquist, K.; Brunow, G.; Lu, F.; Kim, H.; Schatz, P.F.; Marita, J.M.; Hatfield, R.D.; Ralph, S.A.; Christensen, J.H.; et al., Lignins: natural polymers from oxidative coupling of 4-hydroxyphenylpropanoids, *Phytochem Rev* **2004**, *3*, 29-60.
- Weng, J.K.; Li, X.; Bonawitz, N.D.; Chapple, C., Emerging strategies of lignin engineering and degradation for cellulosic biofuel production, *Current Opinion in Biotechnology* **2008**, *19*, 166-172.

3. Wang, W.D.; Yan, L.; Cui, Z.J., Characterization of a microbial consortium capable of degrading lignocellulose, *Bioresour. Technol* **2011**, *120*, 9321–9324.
4. Bugg, T.D.H.; Ahmad, M.; Hardiman, E.M.; Singh, R., The emerging role for bacteria in lignin degradation and bio-product formation, *Current Opinion in Biotechnology* **2011**, *22*, 394–400
5. Barclay, C.D.; Legge, R.L.; Farquha, G.F., Modelling the growth kinetics of *Phanerochaete chrysosporium* in submerged static culture, *Appl. Environ. Microbiol* **1993**, *59*, 1887–1892.
6. Camarero, S.; Sarkar, S.; Ruiz-Duenãs, F.J.; Martnez, M.J.; Martínez, A.T., Description of a versatile peroxidase involved in the natural degradation of lignin that has both manganese peroxidase and lignin peroxidase substrate interaction sites, *J Biol Chem* **1999**, *274*, 10324–10330.
7. Bhat, M.K., Cellulases, Related enzymes in biotechnology, *Biotechnol. Adv* **2000**, *18*, 355–383.
8. Coughlan, M.P., Enzymatic hydrolysis of cellulose: an overview, *Biore-sour. Technol* **1992**, *39*, 107–115.
9. Thurston, C.F., The structure and function of fungal laccases, *Microbiology* **1994**, *140*, 19–26.
10. Pieper, D.H., Aerobic degradation of polychlorinated biphenyls. *Appl Microbiol Biotechnol* **2005**, *67*, 170–191.
11. Adav, S.S.; Ng, C.S.; Arulmani, M.; Sze, S.K., Quantitative iTRAQ secretome analysis of cellulolytic *Thermobifida fusca*, *J Proteome Res* **2010**, *9*, 3016–3024.
12. Brown, M.E.; Chang, M.C.Y., Exploring bacterial lignin degradation, *Current Opinion in Chemical Biology* **2014**, *19*, 1–7.
13. Hamelinck, C.N.; van Hooijdonk, G.; Faaij, A.P.C.; Ethanol from lignocellulosic biomass: techno-economic performance in short-, middle- and long-term, *Biomass Bioenergy* **2005**, *28*, 384–410.
14. Woo, H.L.; Hazen, T.C.; Simmons, B.A.; DeAngelis, K.M., Enzyme activities of aerobic lignocellulolytic bacteria isolated from wet tropical forest soils, *Systematic and Applied Microbiology* **2014**, *37*, 60–67.
15. Gilkes, N.R.; Kilburn, D.G.; Miller, R.C.; Warren, R.A.J., Bacterial cellulases, *Bioresour. Technol* **1991**, *36*, 21–35.
16. Maki, M.; Leung, K.T.; Qin, W., The prospects of cellulase-producing bacteria for the bioconversion of lignocellulosic biomass. *Int. J. Biol. Sci* **2009**, *5*, 500–516.
17. Hatakka, A., Lignin-modifying enzyme from selected white-rot fungi: production and role in lignin degradation. *FEMS Microbiol. Rev* **1994**, *13*, 125–135.
18. Vicuña, R., Bacterial degradation of lignin, *Enzyme Microb. Technol* **1988**, *10*, 646–655.
19. Nilsson, T.; Daniel, G., *Int. Res. Group on Wood Preservation, Working Group 1. Biological Problems.* IRG/WP/ 1235, 1983, 14 pp.
20. Daniel, G.F.; Nilsson, T.; Singh, A.P., *Can. J. Microbiol* **1987**, *33*, 943–946
21. Nilsson, T.; Singh, A.P., *Int. Res. Group on Wood Preservation, Working Group 1. Biological Problems.* IRG/WP/ 1186, 1983, 19 pp.
22. Janshekar, H.; Fiechter, A., *Adv. Biochem. Eng. Biotechnol.* **1983**, *27*, 119–178
23. Kirk, T.K.; Farrell, R., *Annu. Rev. Microbiol* **1987**, *41*, 465–505
24. Burlage, R.S.; Hooper, S.W.; Sayler, G.S., The TOL (pWwo) catabolic plasmid, *Appl Environ Microbiol* **1989**, *55*, 1323–1328.
25. Masai, E.; Katayama, Y.; Nishikawa, S.; Fukuda, M., Characterization of *Sphingomonas paucimobilis* SYK-6 genes involved in degradation of lignin-related compounds, *J Ind Microbiol Biotechnol* **1999**, *23*, 364–373.
26. Zimmermann, W.; Degradation of lignin by bacteria. *Journal of Biotechnology* **1990**, *13*, 119–130
27. Lynd, L.R.; Weimer, P.J.; Van Zyl, W.H.; Pretorius, I.S., Microbial cellulose utilization: fundamentals and biotechnology, *Microbiol. Mol. Biol. Rev* **2002**, *66*, 506–577.
28. Pérez, J.; Muñoz-Dorado, J.; Dela Rubia, T.; Martínez, J., Biodegradation and biological treatments of cellulose, hemicellulose and lignin: an overview, *Int. Microbiol* **2002**, *5*, 53–63.
29. Yang, J.S.; Ni, J.R.; Yuan, H.L.; Wang, E.T.; Biodegradation of three different wood chips by *Pseudomonas* sp. PKE117, *Int. Biodeterior. Biodegrad* **2007**, *60*, 90–95.
30. Das, M.T.; Budhraj, V.; Mishra, M.; Thakur, I.S., Toxicological evaluation of paper mill sewage sediment treated by indigenous dibenzofuran-degrading *Pseudomonas* sp., *Bioresour. Technol* **2012**, *110*, 71–78.
31. Dagley, S., *Adv. Microb. Physiol.* **1971**, *6*, 1–46
32. Chamkha, M.; Garcia, J.L.; Labat, M., Metabolism of cinnamic acids by some Clostridiales and emendation of the descriptions of *Clostridium aerotolerans*, *Clostridium celerecrescens* and *Clostridium xylanolyticum*, *Int. J. Syst. Evol. Microbiol* **2001**, *51*, 2105–2111.

33. Geib, S.M.; Filley, T.R.; Hatcher, P.G.; Hoover, K.; Carlson, J.E.; Jimenez-Gasco, M.M.; Lzumi, A.N.; Sieghter, R.L.; Tien, M., Lignin degradation in wood-feeding insects, *Proc. Natl. Acad. Sci* **2008**, *105*, 12932–12937.
34. Chung, S.Y.; Maeda, M.; Song, E.; Horikoshi, K.; Kudo, T., A Gram-positive polychlorinated biphenyl-degrading bacterium, *Rhodococcus erythropolis* strain TA421, isolated from a termite ecosystem, *Biosci Biotechnol Biochem* **1994**, *58*, 2111-2113.
35. Wenzel, M.; Schönig, I.; Berchtold, M.; Kämpfer, P.; König, H., Aerobic and facultatively anaerobic cellulolytic bacteria from the gut of the termite *Zootermopsis angusticollis*, *J Appl Microbiol* **2002**, *92*, 32-40.
36. Kato, K.; Kozaki, S.; Sakuranaga, M., Degradation of lignin compounds by bacteria from termite guts, *Biotechnol Lett* **1998**, *20*, 459-462.
37. Harazono, K.; Yamashita, N.; Shinzato, N.; Watanabe, Y.; Fukatsu, T.; Kurane, R., Isolation and characterization of aromaticdegrading microorganisms from the gut of the lower termite *Coptotermes formosanus*, *Biosci Biotechnol Biochem* **2003**, *67*, 889-892.
38. Schloss, P.D.; Delalibera, I.; Handelsman, J.; Raffa, K.F., Bacteria associated with the guts of two wood-boring beetles: *Anoplophora glabripennis* and *Saperda vestita*, *Environ Entomol* **2006**, *35*, 625-629.
39. Vasanthakumar, A.; Delalibera, I.; Handelsman, J.; Klepzig, K.D.; Schloss, P.D.; Raffa, K.F., Characterization of gut-associated bacteria in larvae and adults of the southern pine beetle, *Dendroctonus frontalis* Zimmermann, *Environ Entomol* **2006**, *35*, 1710-1717.
40. Delalibera, I.; Vasanthakumar, A.; Burwitz, B.J.; Schloss, P.D.; Klepzig, K.D.; Handelsman, J.; Raffa, K.F., Composition of the bacterial community in the gut of the pine engraver, *Ips pini* colonizing red pine, *Symbiosis* **2007**, *43*, 97-104.
41. Wang, Y.; Liu, Q.; Yan, L.; Gao, Y.; Wang, Y.; Wang, W.; A novel lignin degradation bacterial consortium for efficient pulping, *Bioresource Technology* **2013**, *139*, 113–119
42. Ramachandra, M.; Crawford, D.L.; Hertel, G., Characterization of an extracellular lignin peroxidase of the lignocellulolytic actinomycete *Streptomyces viridosporus*, *Appl. Environ. Microbiol* **1988**, *54*, 3057–3063.
43. Katayama, Y.; Nishikawa, S.; Murayama, A.; Yamasaki, M.; Morohoshi, N.; Haraguchi, T.; The metabolism of biphenyl structures in lignin by the soil bacterium (*Pseudomonas paucimobilis* SYK-6), *FEBS Lett* **1988**, *233*, 129–133.
44. Ahmad, M.; Taylor, C.R.; Pink, D.; Burton, K.; Eastwood, D.; Bending, G.R.; Bugg, D.H., Development of novel assays for lignin degradation: comparative analysis of bacterial and fungal lignin degraders, *Mol. BioSyst* **2010**, *5*, 815–821.
45. Morii, H.; Nakamiya, K.; Kinoshita, S., Isolation of lignin decolouring bacterium, *J. Ferment. Bioeng* **1995**, *80*, 296–299.
46. Raj, A.; Chandra, R.; Krishna Reddy, M.M.; Purohit, H.J.; Atya, K., Biodegradation of kraft lignin by a newly isolated bacterial strain, *Aneurinibacillus aneurinilyticus* from the sludge of a pulp paper mill, *World J. Microbiol. Biotechnol* **2007**, *23*, 793–799.
47. EL-Hanafy, A.A.; Abd-Elsalam, H.E.; Hafez, E.E., Molecular characterization of two native Egyptian ligninolytic bacterial strains, *Res. J. Appl. Sci. Eng* **2008**, *4*(10), 1291–1296.
48. Chandra, R.; Raj, A.; Purohit, H.J.; Kapley, A., Characterisation and optimisation of three potential aerobic bacterial strains for kraft lignin degradation from pulp paper waste, *Chemosphere* **2007**, *67*, 839–846.